

DOI: <https://doi.org/10.65814/ijar.2022.26.1.196>

FUNGI ASSOCIATED WITH *Sesamum indicum* L. SEEDS AND OCHRATOXIGENIC POTENTIAL OF SOME *Aspergillus* AND *Penicillium* ISOLATES IN DUHOK

A. A. M. Saadullah¹F. TH. Ibrahim¹L. F. Hussein¹**Keywords:** OchratoxinA, *Aspergillus*, *penicillium*, sesame seed, ELISAE-mail: asia.saadullah@uod.ac© 2022 Directorate of Agricultural Research, Ministry of Agriculture. This is an-open access article under the CC by Licenses <http://creativecommons.org/licenses/by/4.0>

ABSTRACT

Sesame the oil-based seed crop *Sesamum indicum* L. has been widely planted and consume in Iraq. Unfortunately, sesame seeds are particularly susceptible to different infectious agents especially fungi in the field specially if stored improperly. Using culture and immunological approaches, the current study tried to analyze the ochratoxigenic potential of several *Aspergillus* and *Penicillium* isolates from sesame samples in Duhok city.

The results showed the isolation of 24 fungal species belong to 14 genera with number of sterile mycelia and yeasts, also showed the potency of 17 isolates of *Aspergillus* and *Penicillium* to produce OTA. Ochratoxin A (OTA) were detected in cultures of six species of section nigri (*A.carbonarius* and *A.niger* aggregate), one of *Aspergillus* section circumdati species (*A .ochraceus*) and one species of *Penicillium* (*P.verrucosum*). OTA was found at level from 0.63 to 0.70 ng\ml in *A.carbonarius* isolates; from 0.60 to 0.64 ng\ml in *A.niger* aggreg. isolates; in *P.verrucosum* from 0.52 to 0.61 ng\ml and in *A.ochraceus* isolates 0.23 -0.62 ng\ml while none of both *A.sclerotiorum*, *A.japonicus* isolate produce. According to our findings, the occurrence of ochratoxigenic *Aspergillus* and *Penicillium* isolates in sesame seed samples poses a specific risk to consumer health.

INTRODUCTION

The sesame (*Sesamum indicum* L.) belongs to the family Pedaliaceae and one of the world's most and oldest important oil-seeds. It contains 58 to 44 percent oil, 25 percent protein. 13.5 percent carbohydrates and 5 percent Ash. It also contains lot of dietary fibers and micronutrient including iron, phosphate, calcium and potassium, as well as vitamins E, and niacin, thiamine, tocopherols, lignin, and phyto-sterols (8). Sesame is significant in Iraq due of its several applications.

Fungal genera associated with sesame seeds such as *Alternaria*, *Curvularia*, *Fusarium*, *Helminthosporium*, *Memnoniella*, *Penicillium*, and *Rhizophus* sp. have been observed associated with sesame, causing seed degeneration in soil before

¹ Biology Department, College of Sciences., University of Duhok, Duhok, Iraq

Received: May/2022.

Accepted: Aug. /2022.

germination, seedling mortality, and foliage infection at the adult stage Altaf et al. (8) *Alternaria* sesame is the most damaging disease of sesame, causing little brown patches on leaves ranging from 1 to 8 mm in diameter and reducing seed viability. *Fusarium* sp. causes dark necrotic patches on sesame leaves and yellowing of the foliage is the initial indication of *Fusarium* sp. infection Abarca et al. (2) Seed rot caused by *Aspergillus* and *Fusarium* caused seed rot, which inhibited seed germination. *Fusarium moniliforme* causes root rot and seedling blight by causing brown necrotic lesions on roots and later becoming a seedling invader.

Contamination from fungal agricultural pests, on the other hand, is a big worry. Nowadays, people pay a lot of knowledge to food -safety and pollutants (35, 55). Fungi-created mycotoxin are among the greatest common food pollutants, posing a threat food safety, to public health, and the economies of many countries, particularly developed ones. Moisture, inherent characteristics and long shelf life, nutrients, high water activity and pH are the most important elements in mycotoxin contamination of Food (53). *Aspergillus* fungus, particularly *A. ochraceus* and *A. carbonarius* and occasionally *A. niger* produce ochratoxins OTA.

Sesame (*Sesamum indicum* L.) is a popular tropical oil seed crop sub-tropical and temperate locations around the world, covering approximately 9,398,770 hectares (ha). Sesame is regarded as a nutrient-dense food as a result of its high oil content, which ranges from 44 to 58% proteins and its nutritious, cosmetics qualities and medicinal (29). Sesame seeds have been proven to have anti-oxidant, anticancer, wound-healing, antihypertensive, neuro-protective, anti-inflammatory properties due to the presence of different bioactive chemicals (23, 39).

The majority of Ochratoxin A (OTA) was first discovered in *Aspergillus* section *Circumdati* (*A. ochraceus*) and *Penicillium* species (*P. nordicum* and *P. verrucosum*) Other species such as *P. crustosum*, *P. chrysogenum*, *P. brevicompactum*, *P. oxalicum* and *P. olsonii* have been claimed as ochratoxin A developers (35). However, multiple studies in the recent decade have found that few species of the *Aspergillus* section *nigri* (*A. niger* aggr. and *A. carbonarius*), which are frequent contaminants of fruits like grapes and their derivatives, are ochratoxigenic (3, 22). In *Aspergillus* section *Circumdati* (previously the *Aspergillus ochraceus* group), the follows species can produce OTA: *Aspergillus flocculosus*, *A. cretensis*, *A. roseoglobulosus*, *A. pseudoelegans*, *A. sclerotiorum*, *A. sulphureus* and *Neopitomyces muricatus*. They are the primary cause of OTA contamination in grapes and grapevine products (1, 25). In the Province of Duhok, no investigations on mycotoxins contamination in sesame seeds have been undertaken. As a result, the current investigation attempted to assess the ochratoxigenic ability of various *Aspergillus* and *Penicillium* isolate from sesame samples using cultural and Immunological methods (ELISA) Figure1.

MATERIALS AND METHODS

1 SAMPLES COLLECTION



Figure1: some of the sesame seed samples used in current study.

During August 2021 to January 2022, 100 samples (250 g) of sesame seed were collected in sterile polyethylene bags from different places in Duhok province, north of Iraq. Sesame seeds were transformed to the University of Duhok Mycology Research Laboratory to be tested for fungal contamination, ochratoxin testing, and fungal identification. The seeds were confined to 4°C until they were needed.

2 Isolation and Identification of Fungi

The direct plating method as used to analyze sesame seed samples (034). One hundred seeds were surface sterilized with 2% sodium hypochlorite solution for 2 min. at room temperature, followed by two washes with sterilized distilled water and placement on three culture media using international seed testing association ISTA techniques (10-20 seeds per plate): and used different agar culture (Dichloran rose bengal chloramphenicol Agar medium (DRBC) (Germany, Fluka), malt extract Agar (MEA) medium (France, Biokar Diagnostics), and Potato dextrose agar (PDA) (LAB-M, UK). Plates were incubated for 5-7 days at 28°C. The fungus that grew on the seeds were identified either directly or by sub culturing them on another plates of (PDA) and Czapeck's agar media. The identification of fungal species was done using descriptions from earlier investigations (2, 15, 18, 24, 34, 54).

3 Percentage of Frequency

The following formula was used to calculate the percentage frequency of species isolated.

$$\text{Isolation Frequency \%} = \frac{\text{Number of samples on which a fungus appeared}}{\text{Total number of samples}} \times 100$$

4 Identification of *Aspergillus* and *Penicillium* species

Daily, samples of sesame seed were investigated for sporulation fungi using a stereomicroscope (AmScope, China). For identification, pure colonies were produced on appropriate media. Based on physical and cultural features, the majority of observed taxa were identified to species level. Pure colonies of *Penicillium* and *Aspergillus* were grown on four media for identification (24). The following media were used: Czapeck yeast extract agar (CYA25) (K₂HPO₄)

41 gr, Czapek concentrates 10gr, Powdered Yeast Extract 5 gr, sucrose 30gr, agar 15gr and distilled water 1 liter, incubated for seven days at 25°C, czapeck yeast extract agar (CYA37) incubated for seven days at 37°C, czapeck yeast extract agar with 20% Sucrose incubated for seven days at 25°C (CY20S) K_2HPO_4 1 gr, Czapek concentrates 10gr, Powdered Yeast Extract 5 gr, sucrose 200 gr, agar 15gr and distilled water 1 liter incubated for seven days at 25°C (24, 34). For each culture, two plates of CYA and one plate each of CY20S and MEA were used. Each plate is inoculated in the center and incubated for seven days in the light at 37°C, one CYA is incubated. The others are being incubated at 25 degrees Celsius.

5 Confirmation test for *Aspergillus* and *Penicillium* species identification

5a Ehrlich test

The Ehrlich test was developed by Frisvad et. al (17) to distinguish species of the *Penicillium* subgenus *Penicillium* by detecting alkaloids interacting with Ehrlich reagent (18) using a filter paper method. The Ehrlich reagent is 2 g 4-dimethylamino benzaldehyde in 96 percent ethanol (85 ml) combined with 15 ml 10 N HCl. On the mycelia side of an established colony on CYA (incubated for 5–9 days at 25°C), a four-mm agar plug is cut out, and a piece of wetted filter paper (Whatman No. 1) is placed on the mycelia side. In section Nigri Samson et. al (44) employs this method to classify numerous *Aspergillus* species.

5b Growth on Creatine Sacrose Agar (CREA)

On Creatine Sucrose Agar medium (creatine($1H_2O$), 3g; sucrose, 30g; KCl, 0.5g; $MgSO_4 \cdot 7H_2O$, 0.5g; $FeSO_4 \cdot 7H_2O$, 0.1g; $K_2HPO_4 \cdot 3H_2O$, 1.3g; bromocresol purple, 0.05g; agar, 15.0g and distilled water, 1L), the growth capacities of all *Aspergillus* species in section Nigri isolates were examined CREA is a semi-selective medium that can be used to classify diverse fungal cultures, particularly *Penicillium* species. Formation of acid (changing the medium from purple to yellow) and production of base can be employed as diagnostic features on CREA colonial growth characteristics. CREA can be used to divide all black aspergilli into groups in a semi-selective manner (44). All species were identified using the keys and descriptions provided by the author (2, 18, 24, 43, 45).

6 Determination

The toxigenic potential of fungus in culture media Cabanes et. al (13) and Saito and Machida (43) Saito and Cabanes rapid methods for detecting ochratoxigenic isolates section Nigri based on color change following exposure to ammonia vapor, were used to study the ochratoxigenic potential of all reported *Aspergillus* isolates from section Nigri. Coconut cream agar Petri plates were used to grow the isolates. The medium was prepared according to the instructions of Dyer and McCammon (14), Lin and Dianese (26) Each strain was inoculated in the center of solidified coconut cream agar medium in 9-cm Petri dishes and incubated at 27°C in the dark. After 4 days of incubation, the dishes were flipped upside down and a drop of ammonia solution was placed into the lid of the dish to see the color shift of the colony reversal.

7 Extraction of Ochratoxin A from Fungal culture

The ability of isolates from the *Aspergillus* and *Penicillium* genera to produce OA was tested. The extraction of ochratoxin A was done using the Bragulat et. al. (12) technique. The isolates were cultivated for 7 days on Czapek

yeast autolysate agar (CYA) medium at 25°C. Each colony had three plugs (7 mm diameter) removed from the inner, middle, and outside portions. Plugs were placed in 3-ml vials, 1 ml methanol was added, the vials were shaken, and the vials were incubated at 25°C for 60 minutes. The extracts were centrifuged three times at 4000 rpm for five minutes each time. The supernatant was filtered via a Millipore filter (0.22 μ m) filtration membrane (Millex GP Filter unit Corningwollahill Co Ireland). The extracts were then analyzed using ELISAs (Enzyme linked Immunosorbent Assays) Figure 2.




Figure 2: Green fluorescent a on the back of a colony developed on coconut cream agar (an Ochratoxin A (OTA)-producing strain) (365 nm). Under the same circumstances, a non-OTA producing strain showed no fluorescence (b).

8 Ochratoxin Analysis

The enzyme linked immunosorbent assay was used to produce a quantitative study of OA (ELISA). This was done in accordance with the manufacturer's instructions (Veratox quantitative ochratoxin test-Neogen Corporation-USA). The OTA concentration was estimated using a standard curve derived from OTA standards and expressed in ng/ml.

9 Preservation of the Isolated isolates


Preservation of all fungi isolates was done using culture of PDA in sterilized vial glass and preservation with glycerol and stored at 5°C in a refrigerator.

RESULTS AND DISCUSSION

Fungal species, frequency percentages of genera, and species of fungus isolated during the investigation are shown in Table 1 and figure 3 there were 15 genera and 24 species identified from 100 samples, as well as unidentified yeasts and sterile mycelium. Seven species of *Aspergillus* were found, with the most diversity of all of the retrieved taxa. Three species of black aspergilli were found (*A. carbonarius* 1.3%, *A. japonicas* 0.44% and *A. niger* 3.0%). Other aspergillus species include *A.ochraceous* 1.5%, *A.flavus* 1.0% ,*A.fumigatus* 0.33% and *A.sclerotiorum* 1.0%.In the Kurdistan region, several species were detected in soil and other agricultural items (7, 40) The possibility for contamination of commodities by *Aspergillus* and their toxins is increased by climatic circumstances characterized by humidity and temperature, as well as poor storage techniques 0(36) and Several investigators have noted the abundance of *Aspergillus* in preserved nuts bean (7, 39) figure 4.

Table 1: The genera and species of fungal isolates and the frequency percentages

S.N	Frequency (%) of fungal genera on 7PDA MEA and DRBC 1	
	Fungal genera	(%) Frequency
1	<i>Absidia</i> sp.	2.0
2	<i>Alternaria alternata</i>	0.5
3	<i>Aspergillus carbonarius</i>	1.3
4	<i>A.flavus</i>	1.0
5	<i>A.fumigatus</i>	0.33
6	<i>A.niger</i>	3.0
7	<i>A.ochraceous</i>	1.5
8	<i>A.japonicus</i>	0.44
9	<i>A.sclerotium</i>	1.0
10	<i>Penicillium citrinum</i>	1.5
11	<i>P.expansum</i>	1.0
12	<i>P.verroucosum</i>	0.5
13	<i>Curvularia</i> sp.	1.0
14	<i>Emericella nidulans</i>	0.77
15	<i>Eurotium amstelodami</i>	0.66
16	<i>Fusarium oxysporum</i>	0.33
17	<i>Geotrichum candidum</i>	0.3
18	<i>Monilia</i> sp.	0.28
19	<i>Mucor circinelloides</i>	0.25
20	<i>Rhizopus stolonifera</i>	0.22
21	<i>Sterile mycelium</i>	0.17
22	<i>Trichoderma</i> sp.	0.15
23	<i>Cephalosporium</i> sp.	0.1
24	Yeast non identified	0.1

**Figure 3: Frequency (%) of fungal genera on DRBC.PDA and MEA medium.**




Figure 4a: Growth of *A.niger* and *A.carbonarius* , *Rhizopus stolonifer* and *M.circinelloides* on sesame seed bean on DRBC, b. Growth of *A.niger* and *A.japonicus* and *P.expansum* on sesame seed bean on DRBC medium, c. Growth of *A.niger* and *A.japonicus* and *P.verrocusum* on sesame seed bean on DRBC medium, d. Growth of *A.niger* and *A.carbonarius* and on sesame seed bean on MEA medium, e. Growth of *A.niger* and *A.flavus* and *A.carbonarius* on sesame seed bean on MEA medium, f. Growth of *A.fumigatus* and *E.amstelodami* on sesame seed bean on PDA medium, g Growth of *A.niger* and *P.citrinum* ,*A.flavus* and *A.carbonarius* on sesame seed bean on PDA medium h. Growth of *A.niger* and *A.japonicus*, and *A.ochraceous* on sesame seed bean on MEA.

Penicillium was second in the number of isolated species from seed and was represented by three species. *P.citrinum* was the most common species (1.5 %), followed by *P. expansum* (1%) and *P. verrucosum* (0.5%). *Aspergillus* and *Penicillium* species mainly grow during storage. Two teleomorphic ascomycetes, namely, *Emericella nidulans* and *Eurotium amstelodami* were detected with percentage frequencies 0.77% and 0.66% respectively. Isolates of *E. amstelodami* from turkish dried figs produced Aflatoxin B1 and ochratoxin A when grown on potato- dextrose -broth (PDB) medium0 (47).

Other genera and species include *Absidia* sp., *Alternaria alternate*, *Curvularia* sp., *Fusarium oxysoprum*, *Geotrichum candidum*, *Monilia*, *Mucor circinelloides* ,*Rhizopus stolonifer* ,*Trichoderma* sp *Cephalosporium* sp. with frequency percentages of (2.0, 0.5, 1.0, 0.33, 0.3, 0.28, 0. 25, 0.22, 0.15)% and 0.1% and yeast 1%.

After 7 days at 25°C, the capacity of the fungal isolates to produce OA was tested using solid laboratory medications (CYA). As indicated in 2.6, a fast method of extraction of OA by Bragulat et. al. ⁰(12) was used. Table 3.

Eleven isolates belonged to four species wrer showed ochratoxigenic potential,two pecies from section Nigri (*A.carbonarius*, *A.niger aggr.*), one *Aspergillus* species from section circumdati (*A.ochraceus*), and one species of *Penicillium* were found to contain ochratoxin A. (*P.verrucosum*). For *A.carbonarius*, *A.ochraceus* the proportion of OTA generating isolates examined by ELISA technique was 100 percent, while the percentages of ochratoxigenic potential in *A.niger aggr.* and *P.verrucosum* were 60 percent and 66.6 percent,

respectively (Table 3). Isolates from *A. carbonarius* and, to a lesser extent, *A. niger*, demonstrated the ability to produce ochratoxin A and were isolated often from dried vines naturally contaminated with ochratoxin A in the Kurdistan region (41).

The proportion of ochratoxigenic isolates in the genera *Aspergillus* and *Penicillium* reported from various parts of the world varies considerably based on the number of isolates studied and geographical regions. Several publications have investigated the ochratoxigenic potential of *A. carbonarius* strains (3, 24, 29). All of these research revealed that this species has a constant ability to create OA, with ochratoxigenic potential of *A. carbonarius* isolates ranging from 41.7 to 100%.

Several publications have reported a percentage of ochratoxigenic isolates in *A. niger* ranging from 0.8 to 18.5 percent (3, 09, 037, 50).

Magnoli revealed that 30% of *A. niger* from Brazilian grapes was an OA producer and (28, 51) discovered an abnormally high percentage (43.1%) of ochratoxigenic strains in *A. niger* from Italian grapes.

The uniseriate black *Aspergillus japonicus* showed no ochratoxigenic producing ability (Table 2). This is in line with what is mentioned by literature (11, 24, 28). All the three isolates of *A. ochraceus* were positive for ochratoxin A producing ability. However Serra et. al. (50) and according to the study, 50 percent of *A. ochraceus* strains isolated from Portuguese wine grapes were able to produce OA. Some authors in Argentina, Brazil, and Spain have reported a greater percentage of OA positive isolates among *A. ochraceus*. (20, 28).

Out of 3 strains of *P. verrucosum*, two were positive for OA production. *P. verrucosum* is the major species producing OA in cereals such as wheat and barley in temperate and cold climate Cabafies et. al. (13) however, the fungus was isolated from samples of dried vine grapes.

Table 2: Percentage (%) of ochratoxin A producing strains and range of OTA detected

Fungus (from culture)	Ochratoxin A ng/ml
<i>A. japonicus</i>	0.00
<i>A. carbonarius</i>	0.65
<i>A. carbonarius</i>	0.63
<i>A. carbonarius</i>	0.65
<i>A. ochraceus</i>	0.62
<i>P. verrucosum</i>	0.52
<i>A. niger</i> aggr.	0.64
<i>A. niger</i> aggr.	0.00
<i>A. carbonarius</i>	0.70
<i>P. verrucosum</i>	0.00
<i>A. niger</i> aggr.	0.62
<i>P. verrucosum</i>	0.00
<i>A. sclerotiorum</i>	0.00
<i>A. ochraceus</i>	0.23
<i>A. niger</i>	0.00
<i>A. ochraceus</i>	0.21
<i>A. niger</i> aggr.	0.60

Table 3: List of fungal isolate examined for ochratoxin A by ELISA technique

Species	No. of isolates tested	OA producing isolate (%)	Rangeng/ml.
<i>A.carbonarius</i>	4	100	0.63-0.70
<i>A.japonicus</i>	1	0	N.D
<i>A.niger</i> aggr.	4	75	0.60-0.64
<i>A.ochraceus</i>	2	100	0.23-0.62
<i>A.sclerotiorum</i>	1	0	N.D
<i>P.verrucosum</i>	3	66.6	0.52-0.61

N.D =Not detected.

The possibility for contamination of commodities by *Aspergillus* and their toxins is increased by climatic circumstances characterized by humidity and temperature, as well as poor storage techniques Riba et. al. (36) and several investigators have noted the abundance of *Aspergillus* in preserved nuts bean (10, 39).

CONCLUSION

According to our findings, the occurrence of ochratoxigenic *Aspergillus* and *Penicillium* isolates in sesame seed samples poses a specific risk to consumer health. As a result, such contaminated products should be examined for the presence of fungi. It's crucial to identify the fungi in order to determine out which mycotoxins are present inexpensive culture approaches for detecting ochratoxigenic contamination when restricted resources prevent the use of analytical procedures. To limit the risk of fungal infection, seeds should be treated with a suitable chemical before sowing.

REFERENCES

- 1- Abarca, Ml; F. Accensi; MR. Bragulat; G. Castella and FJ. Cabanes (2003). *Aspergillus carbonarius* as the main source of ochratoxin A contamination in dried vine fruits from the Spanish market. J. of food protection. Mar, 66(3):504-6.<https://www.deepdyve.com/lp/american-society-for-microbiology/>
- 2- Abarca, Ml.; F. Accensi; J. Cano and FJ. Cabañes (2004). Taxonomy and significance of black aspergilli. Antonie van Leeuwenhoek. Aug., 86(1):33-49.<https://www.semanticscholar.org/paper/Taxonomy-and-significance-of...>
- 3- Abarca, Ml; MR. Bragulat; G. Castella and F. Cabanes (1994). Ochratoxin A production by strains of *Aspergillus niger* var. *niger*. Applied and environmental microbiology. Jul., 60(7):2650
[2.https://www.deepdyve.com/lp/american-society-for-microbiology/](https://www.deepdyve.com/lp/american-society-for-microbiology/)
- 4- Abdullah, SK and A.A. Saadullah (2012). Detection of ochratoxigenic potential of *Aspergillus* strains isolated from vineyard soil, fresh grape berries and dried vine fruits by LC-MS/MS technique. on Biological Sci., May8;8:260.https://www.researchgate.net/publication/268925113_Detection_of_
- 5- Abdullah. SK. and AA. Saadullah (2013). Soil mycobiota at grapevine plantations in Duhok, North Iraq. Mesopotamia J. of Agric., 41(1):437-47.
- 6- Abdullah. S.K. and A.A.H. Muhammed (2011). Prevalence of Black Aspergilli in soil at Vineyards in Duhok. Kurdistan Region of Iraq. Proc. 4th Int. Sci. Conf. Salahaddin Univ. Erbil, Oct., 18-20,2011..3,966-969.

- 7- Adebaio. Lo. and SA. Divaolu (2003). Mycology and spoilage of retail cashew nuts. African J. of Bio., 2(10): 369-73; https://www.researchgate.net/publication/284719503_Fungal_contamination_and_mycotoxinproduction_by_Aspergillus_spp_isolated_from_nuts_and_sesame_seeds.
- 8- Altaf, N; SA. Khan; M. Ahmad; R. Asghar; RA. Ahmed; S. Shaheen; M. Zafar and M. Saqib (2004). and Seed borne mycoflora of sesame (*Sesamum indicum* L.) and their effect on germination and seedling. Pakistan J. of Bio. Sci.;7(2):243-5.
- 9- Battilani, P.; P. Giorni and A. Pietri. (2003). Epidemiology of toxin-producing fungi and ochratoxin A occurrence in grape. In Epidemiology of mycotoxin producing fungi (pp. 715-722). Springer, Dordrecht. <https://www.semanticscholar.org/paper/Epidemiology-of-Toxin-Producing>
- 10- Bau, M.; MR. Bragulat; ML. Abarca; S. Minguez and FJ. Cabañes (2005). Ochratoxigenic species from Spanish wine grapes. International J. of Food Microbiology. Feb 1;98(2):125-30.
- 11- Bellí, N.; E. Pardo; S. Marín; G. Farré; AJ. Ramos and V. Sanchis. (2004). Occurrence of ochratoxin A and toxigenic potential of fungal isolates from Spanish grapes. J. of the Sci. of Food and Agric. Apr 30;84(6):541-6. <https://portalrecerca.uab.cat/en/publications/ochratoxigenic-species> 84, PP. 541-546. DOI: 10.4236/as.2014.56054
- 12- Bragulat, MR.; ML. Abarca and FJ. Cabañes (2001). An easy screening method for fungi producing ochratoxin A in pure culture. International Journal of food microbiology. Dec 30;71(2-3):139-44. <https://portalrecerca.uab.cat/en/publications/an-easy-screening-method>.
- 13- Cabañes FJ.; MR. Bragulat and G. Castellá (2010). Ochratoxin A producing species in the genus *Penicillium*. Toxins. May; 2(5):1111-20. Toxins 2, PP.1111-1127. <https://pubmed.ncbi.nlm.nih.gov/10419279>
- 14- Dyer SK. and S. McCammon (1994). Detection of toxigenic isolates of *Aspergillus flavus* and related species on coconut cream agar. Journal of Applied Bacteriology. Jan; 76(1):75-8. <https://www.semanticscholar.org/paper/Detection-of-toxigenic-isolates>
- 15- Ellis MB. (1976). More Dematiaceous Hyphomycetes Commonwealth Mycological Institute, Kew. Surrey, England. [https://www.scirp.org/\(S\(vtj3fa45qm1ean45vvffcz55\)\)/reference/](https://www.scirp.org/(S(vtj3fa45qm1ean45vvffcz55))/reference/).
- 16- Fente CA.; JJ. Ordaz; BI. Vazquez; CM. Franco and A. Cepeda (2001). New additive for culture media for rapid identification of aflatoxin-producing *Aspergillus* strains. Applied and environmental microbiology. Oct., 1:67(10):4858-62.
- 17- Frisvad JC.; JM. Frank; JA. Houbbraken; AF. Kuijpers and RA. Samson (2004). New ochratoxin A producing species of *Aspergillus* section Circumdati. Stud. Mycol. Jan., 1;50(1):23-44. [https://www.scirp.org/\(S\(lz5mqp453edsnp55rrgjt55\)\)/reference](https://www.scirp.org/(S(lz5mqp453edsnp55rrgjt55))/reference)
- 18- Frisvad JC. and RA. Samson (2004). Polyphasic taxonomy of *Penicillium* subgenus *Penicillium*. A guide to identification of food and air-borne terverticillate *Penicillia* and their mycotoxins. Studies in mycology. Jan 1;49(1):174. [https://www.scirp.org/\(S\(lz5mqp453edsnp55rrgjt55\)\)/reference](https://www.scirp.org/(S(lz5mqp453edsnp55rrgjt55))/reference).
- 19- Frisvad, JC. and U. Thrane (2000). Mycotoxin production by common filamentous fungi. In Introduction to food-and airborne fungi (Eds.: RA Samson, ES Hoekstra, JC Frisvad and O. Filtenborg) (pp. 321-331). Centraalbureau voor Schimmelcultures. [https://www.Scirp.org/\(S\(lz5mqp453edsnp55rrgjt55\)\)/reference](https://www.Scirp.org/(S(lz5mqp453edsnp55rrgjt55))/reference).

- 20- Hakobyan, L.; K. Grigoryan and A. Kirakosyan (2010). Contamination of raisin by filamentous fungi–potential producers of ochratoxin A. Potravinárstvo (Slovak Republic) <https://www.semanticscholar.org/paper/CONTAMINATION-OF-RAISIN-BY>
- 21- Heenan, CN; KJ. Shaw and JI. Pitt. Ochratoxin A production by *Aspergillus carbonarius* and *A. niger* isolates and detection using coconut cream agar. [https://www.scirp.org/\(S\(lz5mqp453edsnp55rrgjct55\)\)/reference/](https://www.scirp.org/(S(lz5mqp453edsnp55rrgjct55))/reference/).
- 22- Hocking, AD; P. Varelis; JI. Pitt; SF. Cameron and SI. Leong (2003). Occurrence of ochratoxin A in Australian wine. Australian J. of Grape and Wine Res. Apr. 9(1):72-8. <https://www.onlinelibrary.wiley.com/doi/abs/10.1111/j.1755-0238.2003.tb00234.x>.
- 23- Khan, MM.; T. Ishrat; A. Ahmad; MN. Hoda; MB. Khan; G. Khuwaja; P. Srivastava; SS. Raza; F. Islam and S. Ahmad (2010). Sesamin attenuates behavioral, biochemical and histological alterations induced by reversible middle cerebral artery occlusion in the rats. Chemico-Biological Interactions. Jan., 5;183(1):255-63.<https://doi.org/10.1016/j.cbi.2009.10.003>.
- 24- Klich MA. (2002). Identification of common *Aspergillus* species. Centraalbureau voor schimmelcultures. DOI: <https://doi.org/10.1017/S0953756203237783>
- 25- Leong SI.; AD. Hocking and ES. Scott (2007). *Aspergillus* species producing ochratoxin A: isolation from vineyard soils and infection of Semillon bunches in Australia. J. of Applied Microbiology. Jan;102(1):124-33.<https://sfamjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365>.
- 26- Lin MT. and JC. Dianese (1976). A coconut-agar medium for rapid detection of aflatoxin production by *Aspergillus* spp. Phytopathology. Dec., 66(12):1466-9.<https://www.scirp.org/reference/ReferencesPapers.aspx?ReferenceID=1924765>.
- 27- Lund F. Differentiating (1995). *Penicillium* species by detection of indole metabolites using a filter paper method. Letters in Applied Microbiology. Apr., 20(4):228 31.<https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1472-765X.1995.tb00434>.
- 28- Magnoli C.; M. Violante; M. Combina; G. Palacio and A. Dalcero (2003). Mycoflora and ochratoxin-producing strains of *Aspergillus* section *Nigri* in wine grapes in Argentina. Letters in Applied Microbiology. Aug; 37(2):179-84.
- 29- Morris JB. (2009). Characterization of sesame (*Sesamum indicum* L.) germplasm regenerated in Georgia, USA. Genetic Resources and Crop Evolution. Nov., 56(7): 925-36.DOI: 10.4236/as.2014.56054.
- 30- Nakano D.; C. Itoh; F. Ishii; H. Kawanishi; M. Takaoka; Y. Kiso; N. Tsuruoka; Tanaka T. and Y. Matsumura (2003). Effects of sesamin on aortic oxidative stress and endothelial dysfunction in deoxycorticosterone acetate-salt hypertensive rats. Biological and pharmaceutical bulletin.;26(12):1701-5. <https://doi.org/10.1248/bpb.26.1701>.
- 31- Namiki M. (2007). Nutraceutical functions of sesame: a review. Critical reviews in food science and nutrition. Sep., 27;47(7):651-73.<https://www.semanticscholar.org/paper/Nutraceutical-Functions-of-Sesame>.

- 32- Namiki, M. (2007). Nutraceutical functions of sesame: a review. Critical reviews in food science and nutrition. Sep., 27;47(7):651-73. View PDF. CrossRefView Record in Scopus Google Scholar.
- 33- Paterson, RR.; A. Venâncio and N. Lima (2004). Solutions to *Penicillium* taxonomy crucial to mycotoxin research and health. Research in Microbiology. Sep., 1;155(7):507-13.
- 34- Pitt, JI. and AD. Hocking. Fungi and Food Spoilage vol 2 (London, UK: Blackie Academic and Professional).
<https://www.scirp.org/reference/ReferencesPapers.aspx>
- 35- Rapa, M.; S. Ciano; L. Gobbi; R. Ruggieri and G. Vinci (2021). Quality and safety evaluation of new tomato cultivars. Italian J. of Food Sci. Apr., 7;33(2):35-45.
[10.15586/ijfs.v33i2.1921](https://doi.org/10.15586/ijfs.v33i2.1921). <https://doi.org/10.15586/ijfs.v33i2.1921>
- 36- Riba, A.; N. Bouras; S. Mokrane; F. Mathieu; A. Lebrih and N. Sabaou (2010). *Aspergillus* section Flavi and aflatoxins in Algerian wheat and derived products. Food and Chemical Toxicology. Oct., 1;48(10):2772-7. https://www.researchgate.net/publication/284719503_Fungal_contamination_andmycotoxin_production_by_Aspergillus_spp_isolated_from_nuts_and_sesame_seeds [accessed Mar 17 2022].
- 37- Romero, SM.; RM. Comerio; G. Larumbe; A. Ritieni; G. Vaamonde and VF. Pinto (2005). Toxigenic fungi isolated from dried vine fruits in Argentina. International J. of food microbiology. Sep., 25;104(1):43-9.<https://pubmed.ncbi.nlm.nih.gov/16054260>
- 38- Rosa, CD.; V. Palacios; M. Combina; ME. Fraga; AD. Rekson; CE. Magnoli and AM. Dalcero (2002). Potential ochratoxin A producers from wine grapes in Argentina and Brazil. Food Additives & Contaminants. Apr., 1;19(4):408
[14.https://www.tandfonline.com/doi/abs/10.1080/02652030110092748](https://www.tandfonline.com/doi/abs/10.1080/02652030110092748)
- 39- Rostami, R; K. Nadafi; A. Aghamohammadi; SH. Najafi; DM. Fazlzadeh and M. Davil (2009). Survey of peanut fungal contamination and its relationship with ambient conditions in the Bazar of Zanjan. From https://www.researchgate.net/publication/284719503_Fungal_contamination_and_mycotoxin_production_by_Aspergillus_spp_isolated_from_nuts_and_sesame_seeds .
- 40- Saadullah, A. A. and SK. Abdullah (2018). Detection of ochratoxigenic potential in some *Aspergillus* and *Penicillium* isolates from vineyard soil, fresh and dried grapes by ELISA. Oct., 28:1-7. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6524999>.
- 41- Saadullah, A. A. and SK. Abdullah (2012). New Records Of Ochratoxin a Producing Species Of *Aspergillus* Contaminated Raisins From Duhok, Kurdistan Region, Iraq. J. Univ. Duhok., 15(1):377-84. https://www.researchgate.net/publication/268925240Saadullah_AAM
- 42- Sage, L.; S. Krivobok; É. Delbos; F. Seigle-Murandi and EE. Creppy (2002). Fungal flora and ochratoxin A production in grapes and musts from France. J. of Agric. and Food Chemistry. Feb., 27;50(5):1306-11.
- 43- Saito, M. and S. A. Machida (1999). rapid identification method for aflatoxin-producing strains of *Aspergillus flavus* and *A. parasiticus* by ammonia vapor. Mycoscience. Apr 1;40(2):205-8. [https://www.scirp.org/\(S\(i43dyn45teexjx455qlt3d2q\)\)/reference/](https://www.scirp.org/(S(i43dyn45teexjx455qlt3d2q))/reference/).
- 44- Samson, RA.; JA. Houbraken; AF. Kuijpers; JM. Frank and JC. Frisvad (2004). New ochratoxin A or sclerotium producing species in *Aspergillus* section Nigri. Stud Mycol. Jan., 1;50(1):45-56. .

- 45- Samson. RA.; P. Noonim; M. Meier; JA. Houbraken; JC. Frisvad and J.Varga (2007). Diagnostic tools to identify black aspergilli. *Studies in mycology*. Ser 1:59(1):129-45. <https://pure.knaw.nl/portal/en/publications/diagnostic-tools-to>.
- 46- Samson. RA. and J. Varga (2007). editors. *Aspergillus systematics in the genomic era*. Utrecht: CBS Fungal Biodiversity Centre; Apr 12.https://www.aspergillus.org.uk/slide_presentations/ochratoxin-a-in.
- 47- Senyuva, HZ.; J. Gilbert; SÜ. Ozcan and U. Ulken (2005). Survey for co-occurrence of ochratoxin A and aflatoxin B1 in dried figs in Turkey by using a single laboratory-validated alkaline extraction method for ochratoxin A. *Journal of Food Protection*. Jul. 68(7):1512-5.
- 48- Serra, R; L. Abrunhosa; Z. Kozakiewicz and A. Venâncio (2003). Black *Aspergillus* species as ochratoxin A producers in Portuguese wine grapes. *International journal of food microbiology*. Nov., 15;88(1):63-8. <https://pubmed.ncbi.nlm.nih.gov/14527786>.
- 49- Serra, R.; A. Braga and A. Venâncio (2005). Mycotoxin-producing and other fungi isolated from grapes for wine production, with particular emphasis on ochratoxin A. *Res. in Microbiology*. May, 1;156(4):515-21.<https://www.semanticscholar.org/paper/Solutions-to-Penicillium-taxonomy>.
- 50- Serra, R.; C. Mendonça and A.Venâncio. Ochratoxin (2006). A occurrence and formation in Portuguese wine grapes at various stages of maturation. *International J. of Food Microbiology*. Sep., 1;111:S35-9.
- 51- Téren, J.; J.Varga; Z.Hamari; E.Rinyu and F.Kevei (1996). Immunochemical detection of ochratoxin A in black *Aspergillus* strains. *Mycopathologia*. Jun,134(3):171-6.<https://link.springer.com/article/10.1007/s11046-006-0055-4>
- 52- Tjamos, SE.; PP. Antoniou; A. Kazantzidou; DF. Antonopoulos; I. Papageorgiou and EC. Tjamos (2004). *Aspergillus niger* and *Aspergillus carbonarius* in Corinth raisin and wine-producing vineyards in Greece: population composition, ochratoxin A production and chemical control. *J. of Phytopathology* Apr., 152(4):250-5.<https://www.semanticscholar.org/paper/Aspergillus-spp.-distribution>
- 53- Wang, YJ.; JY. Nie; YA. Zhen; LI ZX; Y. CHENG and S. Farooq (2018). Multi-mycotoxin exposure and risk assessments for Chinese consumption of nuts and dried fruits. *J. of Integrative Agric*. Jul., 1;17(7):1676-90.10.1016/S2095-3119(18)61966-5. [https://doi.org/10.1016/S2095-3119\(18\)61966-5](https://doi.org/10.1016/S2095-3119(18)61966-5) www.sciencedirect.com/reference/283289 59.
- 54- Watanabe T. (2002). *Pictorial atlas of soil and seed fungi: morphologies of cultured fungi and key to species*. CRC press; Apr 18.
- 55- Xinyu Z.; Yi W; H. Xue; Y. Zixi; Y. Weiqiong and L. Zhaolin (2020). Characterization of volatile compounds in five blueberry varieties using purge and trap coupled to gas chromatography-mass spectrometry. *Italian J. of Food Sci*. May, 11:32(2).10.14674/IJFS-1675. <https://doi.org/10.1016/j.lwt.2020.110573>

DOI: <https://doi.org/10.65814/ijar.2022.26.1.196>

الفطريات المرافقة لبذور السمسم وقابلية بعض عزلات من الفطرين اسبرجلس وبنسيليوم على انتاج الاوكراتوكسين في محافظة دهوك

آسيا عبد الحميد محمد سعدالله¹ فنا ذكري ابراهيم¹ لينا فرهاد حسين¹

الكلمات الدالة: اوكراتوكسين A، اسبرجلس، بنسيليوم، بذور السمسم. تقنية الاليزا

E-mail: asia.saadullah@uod.ac

© 2022 Directorate of Agricultural Research, Ministry of Agriculture. This is an-open access article under the CC by Licenses <http://creativecommons.org/licenses/by/4.0>



الملخص

محصول السمسم من المحاصيل الزيتية المهمة في العراق، يتعرض المحصول للإصابة بالعديد من المسببات المرضية خاصة الفطريات المرافقة للبذور. تضمنت الدراسة عزل وتشخيص الفطريات المرافقة لبذور السمسم في محافظة دهوك والتحري عن قابلية بعض عزلات الفطرين اسبرجلس وبنسيليوم على إنتاج السم الفطري اوكراتوكسين باستخدام الطرق الزراعية والمناعية. أظهرت النتائج عزل 24 نوعاً تعود الى 14 جنساً فضلاً عن عدد من الفطريات العقيمة والخمائر، أظهرت نتائج قابلية 17 عزلة من الفطرين اسبرجلس و بنسيليوم لإنتاج السم الفطري اوكراتوكسين ستة انواع من مجموعة *nigri* (*A. carbonarius* and *A. niger* aggreg.) ونوع واحد من مجموعة *circumdati* (*A. ochraceus*) ونوع واحد من جنس بنسيليوم (*P. verrucosum*). اوكراتوكسين A وجدت بمعدل (0.63-0.70 ng/ml) في *A. carbonarius* و بمعدل (0.60-0.64 ng/ml) في *A. niger* aggreg. وجدت بمعدل (0.52-0.61 ng/ml) في *P. verrucosum* ; وبمعدل (0.23-0.62 ng/ml) في *A. ochraceus* بينما لم يفرز كل من *A. japonicus* و *A. sclerotiorum*. بخصوص النتائج المكتشفة فإن وجود الفطريات اسبرجلس وبنسيليوم المنتجة للسم الفطري اوكراتوكسين في بذور السمسم يمثل خطراً مهدداً لصحة المستهلك.

¹كلية العلوم، جامعة دهوك، دهوك، العراق.

تاريخ تسلم البحث: آيار/2022.

تاريخ قبول البحث: آب/2022.